


# Spinal Curve and Other Back Deformities


## CHAPTER

# 20

The backbone, or spine, is a chain of bones called vertebrae that connect the head to the hipbone. Separating each of the vertebrae is a small cushion called a disk. The backbone holds the body and head upright. It also encloses, in its hollow center, the spinal cord or trunk line of nerves connecting the brain to all parts of the body (see p. 35).



### SHAPE OF THE BACKBONE



The backbone (and back) is usually straight when seen from the back, and has 4 curves when seen from the side.

### Sideways curve (scoliosis—S-shaped curve)




### Rounded back (kyphosis)



### Swayback (lordosis)




### Sharp bend or bump in spine (tuberculosis of the backbone)




Of these, scoliosis or a sideways curve is the most common serious condition. Often, however, rounded and/or swayback are seen together with scoliosis.

### NON-FIXED AND FIXED SPINAL CURVES

With a non-fixed or "functional" curve there is no deformity of the vertebrae. This usually happens when the body tries to stand straight even though the hips tilt or there is other unevenness not in the spine.



Fixed or "structural" curves are deformities in the bones of the back themselves.



**Note:** In some cases, with time a non-fixed curve may gradually become fixed.

## CAUSES OF SPINAL CURVE (SCOLIOSIS)

Most scoliosis (about 80%) occurs in otherwise healthy children for no known reason. Sometimes it occurs in several members of the same family, so there may be a hereditary (familial) factor. Although about 1 of every 10 persons has some scoliosis (if looked for), only about 1 in 400 has enough of a curve to cause disability. Curves of unknown cause are often first seen—and progress quickly—in children from 10 to 16 years old, during the period of rapid growth.


Known causes of fixed scoliosis range from infection to tumor to rare disease. When possible, consult a doctor with experience in these conditions.

Some children are born with fixed scoliosis, or develop it in early childhood, because of disabilities in the spine itself.

Sometimes one or more vertebrae are only partly formed and cause the spine to bend to one side.



Sometimes 2 or more vertebrae remain attached or “fused” on one side. They can only grow on the unfused side, causing an increasing curve.




Non-fixed scoliosis always results secondary to other disabilities, such as uneven paralysis of the back muscles, or a hip tilt (often due to a shorter leg). Spinal curve often develops in children with polio, cerebral palsy, muscular dystrophy, spina bifida, spinal cord injury, arthritis, and dislocated hip. Be sure to examine all children with these disabilities for spinal curve. With time, non-fixed curves may gradually become fixed.

These conditions can be identified only by X-rays.

### Examining for spinal curve

This is discussed in the chapter on physical examination (Chapter 4).



The rib hump is formed because where the spine is curved, the vertebrae also are twisted to one side.

Look along the line of the back with the child bent over.




### POSSIBLE SIGNS OF EARLY SCOLIOSIS


1. One shoulder higher than other.
2. One shoulder blade sticks out more.
3. One hip is higher or sticks out more than the other.
4. One arm hangs closer to body than the other.
5. Sway back.
6. Rounded shoulders or "hump back."
7. A larger crease at one side of waist than other
8. Child stands sagging or leaning to one side.
9. Rib hump (when child leans forward).
10. A hump near the waist (when leaning forward).
11. A brother, sister, parent, or close relative with scoliosis.

**CHECK FOR:**

one  
shoulder  
lower  
than the  
other



To see the curve better, mark the tip of each vertebra. The actual spinal curve is greater than the curve you have marked.



overhead  
view of  
vertebrae



tips

When you examine for scoliosis, also check to see if the curve can be straightened (non-fixed), or cannot be straightened (fixed).




If untreated, scoliosis may put pressure on a child's internal organs and affect their ability to breathe.

## What to do

This will depend on:

- how severe the curve is.
- if it is getting worse—and if so, how quickly.
- whether the curve is fixed.
- the age of the child.


In a child who does not stand, check spinal curve while he is sitting. If one side of his butt is weaker and smaller, it may cause a hip tilt.



Put a book or board under the weaker butt, and see if this straightens his spine. If so, a cushion raised on one side may help him sit straighter.

How severe the curve is and whether it is getting worse can be best measured by X-rays.


1. For each curve, pick the 2 vertebrae that tilt most in relation to each other.
2. Draw lines level with the top of each vertebra.



3. Measure and record the angle of the spinal curve.
4. Regularly record the curve and notice any changes.

JUAN'S SPINE CURVE	
MAY 86	
JULY 86	
SEPT 86	
NOV 86	
JAN 87	
MAR 87	


Because X-rays are expensive and often hard to get, you can get some idea of whether the curve is getting worse by measuring the angle of the rib hump.



Have the child stand or sit as straight as possible, while he bends forward.

### A homemade level for measuring rib hump angle

Fill the tube with colored water so that the water reaches the top edge of the board at both ends when level.




When measuring the rib hump, level the top by making sure the water is at the top edge of the board at both ends.

If the rib hump angle stays about the same month after month, the curve is probably not getting worse. Keep checking it every few months. If the rib hump angle increases steadily, the curve is getting worse. X-rays should be taken and a decision made about what to do.

Non-fixed curves that are not getting worse should usually be treated only by doing something about the underlying condition.

For example, if the child's spinal curve is not fixed and comes from a hip tilt due to unequal leg length:



This child was developing a spinal curve due to hip tilt and short leg.




Village rehabilitation workers put a lift on his sandal.



This corrected his spinal curve and lop-sided posture.


Body jackets or bracing for a non-fixed curve usually do not help to correct the curve or even to prevent its getting worse. However, for a child with a curve so severe that it makes sitting or walking difficult, a body jacket or corset may help.



Instructions for making body jackets are on p. 558.


Spinal curves under  $20^\circ$  (fixed or non-fixed) usually need only to be watched and measured every few months to see if they are getting worse.

Some experts say that exercises to strengthen the back muscles, like this, help correct and slow down the curving of the spine. Other experts say it does no good. (We do not know.)





Spinal curves over  $20^\circ$ , if they are fixed and getting worse, may get worse less quickly with a brace

A brace like this (the Milwaukee brace) is often used. It works because it is so uncomfortable that the child must stretch his body as straight as possible to reduce the discomfort.



A plastic Boston brace like this is more comfortable, can be completely hidden under the clothes, and probably does as much good.



## SURGERY


For spinal curves over  $50^\circ$  which are quickly getting worse, surgery may be needed. Surgery fuses (joins together) the most affected vertebrae. Usually it only partly straightens the spine. Except for very severe curves, surgery should be avoided in children under 12 years old because the fused part of the spine will not grow any more.

If the curve of the spine is less than  $40^\circ$  by the time the child stops growing, usually it will not progress further. If the curve is over  $50^\circ$ , it is likely to keep getting worse even after the child stops growing, and surgery is often recommended.

However, spinal fusion surgery is very costly and requires an orthopedic surgeon specially trained in this operation. It can also be very hard on the child and family. When surgery cannot be obtained, a body jacket or brace should perhaps be used to help slow down the curve's progress. When a curve becomes too severe, there is no longer enough room in the chest for the lungs and heart to work well, and the child may get pneumonia and die.

## EXERCISES FOR ROUNDED BACK AND SWAYBACK


Children with rounded back may benefit from exercises to help straighten it, like this.



These exercises are explained in the Exercise Sheet #5, p. 387.

The child should also be encouraged to sit and stand as straight as possible, with the shoulders back.

Children with swayback may benefit from exercises to strengthen the stomach muscles, like this,



or by exercises to correct hip contractures (if the child has them). See Exercise Sheet #3, p. 385.